



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Five hundred and ninety-second Meeting.

March 10, 1868. — ADJOURNED STATUTE MEETING.

The PRESIDENT in the chair.

The President called the attention of the Academy to the recent decease of Sir David Brewster of the Foreign Honorary Members, and of Hon. Daniel Lord, of New York, of the Associate Fellows.

Five hundred and ninety-third Meeting.

April 14, 1868. — MONTHLY MEETING.

The PRESIDENT in the chair.

The President called the attention of the Academy to the recent decease of Dr. Samuel L. Dana, of the Resident Fellows, and of Professor William Smyth, of the Associate Fellows.

The following paper was presented : —

*Dispersion of a Ray of Light refracted at any number of Plane Surfaces.** By EDWARD C. PICKERING.

LET $a_1 a_2 a_3$, &c., be the angles included between the surfaces, $n_1 n_2 n_3$ their indices of refraction, $i_1 i_2 i_3$ the angles of incidence, $r_1 r_2 r_3$ the angle of refraction; $\sin i_1 = n_1 \sin r_1$ and in general

$$\sin i_m = r_m \sin r_m \quad (1)$$

$$\text{also} \quad i_m = a_{m-1} + r_{m-1} \quad (2)$$

As the dispersion of any portion of the spectrum is always proportional to the angular divergence of two rays of nearly equal refrangibility, if we vary $n_1 n_2$, &c., dr_m will measure the dispersion. Differentiating (1)

$$\cos i_m di_m = r_m \cos r_m dr_m + \sin r_m dn_m$$

* Since presenting this communication to the Academy, I have learned that a portion of this subject was studied by Sir David Brewster in 1812. I have, therefore, modified my paper, omitting what was not new, except when necessary to preserve the context.